

The Division of Plasma Physics of the American Physical Society is pleased to announce its Distinguished Lecturers for 2012–2013. This Program is intended to share with the larger scientific community the exciting recent advances in plasma physics.

Under the Plasma Physics Travel Grant Program funded by the U.S. Department of Energy, the Lecturers are available for talks at U.S. colleges and universities for the academic year 2012–2013. Their travel expenses will be supported by the grant. No travel-related expenditure is expected from the lecture-hosting institution.

The Lecturers may be invited by contacting them directly.

Additional information about the Plasma Travel Grant Program can be obtained from the Chair of the DPP Education and Outreach Committee:

Paul Miller

Department of Physics

West Virginia University

phone: 304-293-4939

fax: 304-293-5732

email: paul.miller@mail.wvu.edu

Sizing Up Plasmas using Dimensionless Parameters

Craig C. Petty, DIII-D Fusion Science Group

General Atomics, San Diego, CA

petty@fusion.gat.com

Fusion energy, an attractive long-range option for the U.S. clean-energy portfolio, is created when light atomic nuclei are fused together by heating ionized gas (called plasma) to a temperature of hundreds of millions of degrees. Such plasma can be held in place by strong magnetic fields, and while a great deal of progress has been made in understanding this complex physical system using first principle models, dimensionless analysis has also proven to be a valuable tool. This talk reviews what the scaling of phenomena with dimensionless parameters has taught us about the physics of magnetically confined fusion plasmas, and about the extrapolation of present-day experiments to future burning-plasma devices. First, the basic principles of dimensional analysis are introduced by discussing the Buckingham Π theorem and Connor-Taylor scale invariance. Second, the cornerstone principle of similarity, which prescribes that plasmas with the same dimensionless parameters but different physical size exhibit the same physical behavior, is demonstrated. Next, experiments using toroidally configured plasma (i.e., tokamaks) are described that measure the dependence of plasma turbulence and diffusion on dimensionless parameters associated with the normalized plasma size, pressure, and collision frequency. These studies show that plasma turbulence consists of mainly electrostatic fluctuations with characteristic cross-magnetic-field wavelengths on the order of the ion gyroradius (i.e. "micro-scale" turbulence), in general agreement with state-of-the-art drift-wave models. Finally, a modified "wind tunnel" approach to extrapolating dimensionless plasma behavior is discussed that points to a favorable path for increasing the fusion performance in future devices such as the international ITER burning-plasma experiment presently under construction.

Taming turbulence in plasmas: from magnetic fusion energy to black hole accretion disks

Troy Carter, Department of Physics and Astronomy
University of California, Los Angeles, CA
tcarter@physics.ucla.edu

Plasmas in the laboratory and in astrophysical settings vary widely in parameters (e.g. temperature and density) but have one thing in common: they are plagued by instabilities. Instabilities and associated turbulence are detrimental in laboratory plasmas for fusion energy research, causing heat, particles and momentum to "leak" across the confining magnetic field. In astrophysical plasmas like accretion disks, turbulence is essential to explain observed rates of momentum transport and accretion. I will talk about instabilities and turbulence in magnetized plasmas and their relevance to achieving magnetic confinement fusion in the laboratory and understanding processes in astrophysical plasmas.

Novel Physics with Non-neutral Plasmas

Daniel Dubin, Department of Physics
University of California, San Diego, CA
dhdubin@ucsd.edu

Nonneutral plasmas consisting of particles with only a single sign of charge (e.g., pure-electron, pure-positron, or pure-ion plasmas) can be confined for *days or even weeks* using the static electric and magnetic fields of a Penning trap. These novel plasma systems can access regimes unavailable to quasi-neutral plasmas. For instance, they have been cooled to cryogenic temperatures without recombination, and can even form non-neutral *liquid or crystalline* states. In this talk I will review a few aspects of the broadly varied physics accessible to these long-lasting trapped non-neutral plasmas, including applications to precision spectroscopy, quantum computing, fluid dynamics, fusion science, plasma astrophysics, and ongoing efforts to create trapped anti-hydrogen atoms.

Plasmas in Space: From the Surface of the Sun to the Orbit of the Earth

Steven Spangler, Department of Physics and Astronomy
University of Iowa, Iowa City, IA
Steven-spangler@uiowa.edu

Plasma is gas in which a portion of the atoms or molecules are ionized, and thus consist of ions and electrons as well as neutral particles. Plasmas display a richer variety of phenomena than neutral gases. Plasma physics is of importance to astronomy and astrophysics because many astronomical objects are made of plasma. In this talk, I will describe our understanding of the plasma state of the solar atmosphere and interplanetary space (also referred to as the "solar wind") between the Sun and the Earth. The solar atmosphere and interplanetary space are of interest for two reasons. First, they are important astronomical objects and media in their own right. Second, the Sun and interplanetary medium can be measured with a level of detail and precision unattainable for other astronomical objects. In the case of the interplanetary medium, we have the opportunity to make direct "ground truth" measurements of plasma parameters such as density, temperature, and all three vector components of the magnetic field. These more detailed measurements provide a clear view of physical processes which also occur elsewhere in astronomy. I will emphasize those aspects of solar and interplanetary physics which I find most intriguing, and potentially most important in a broader astronomical and astrophysical context. These topics include remote magnetometry of the solar corona (the highest altitude in the solar atmosphere), the attempt to discover the mechanisms responsible for heating the solar corona to temperatures as high as 2 million Kelvin, and the properties of turbulence in the solar atmosphere and interplanetary space. I will conclude by showing how knowledge gained through study of the

Sun and interplanetary medium can contribute to our understanding of much more remote astronomical objects in the Milky Way galaxy and beyond.

Self-Organization - Nature's Intelligent Design

Clint Sprott, Department of Physics

University of Wisconsin, Madison, WI

sprott@physics.wisc.edu

Complex patterns are common throughout nature, from the distribution of the galaxies in the Universe to the organization of neurons in the human brain. It is generally assumed that such complex structure must have a complex cause, but it may be that the patterns spontaneously arise through the repeated application of simple rules. This talk will provide examples of self-organization in nature and will describe six simple computer models that can replicate the features of these patterns. The models typically produce fractal spatial structure and chaotic temporal dynamics characterized by power laws and unpredictability, even when the models are simple and purely deterministic. The work has application to fields as diverse as physics, ecology, political science, economics, sociology, and art.